arXiv:1402.1645v1 [nucl-ex] 7 Feb 2014

^{11}Be(βp), a quasi-free neutron decay?

K. Riisager[a,*], O. Forstner[b,c], M.J.G. Borge[d,e], J.A. Briz[e], M. Carmona-Gallardo[e], L.M. Fraile[f], H.O.U. Fynbo[a], T. Giles[g], A. Gottberg[e,g], A. Heinz[h], J.G. Johansen[a,1], B. Jonson[h], J. Kurcewicz[d], M.V. Lund[a], T. Nilsson[h], G. Nyman[h], E. Rapisarda[d], P. Steier[b], O. Tengblad[e], R. Thies[h], S.R. Winkler[b]

[a] *Department of Physics and Astronomy, Aarhus University, DK–8000, Aarhus C, Denmark*
[b] *Faculty of Physics, University of Vienna, Währinger Strasse 17, A–1090 Wien, Austria*
[c] *Stefan-Meyer-Institut für subatomare Physik, Austrian Academy of Sciences, A–1090 Wien, Austria*
[d] *ISOLDE, PH Department, CERN, CH–1211 Geneve 23, Switzerland*
[e] *Instituto de Estructura de la Materia, CSIC, E-28006 Madrid, Spain*
[f] *Grupo de Física Nuclear, Universidad Complutense de Madrid, CEI Moncloa, E-28040 Madrid, Spain*
[g] *EN Department, CERN, CH–1211 Geneve 23, Switzerland*
[h] *Fundamental Fysik, Chalmers Tekniska Högskola, SE–41296 Göteborg, Sweden*

Abstract

We have observed β^--delayed proton emission from the neutron-rich nucleus ^{11}Be by analysing a sample collected at the ISOLDE facility at CERN with accelerator mass spectrometry (AMS). With a branching ratio of $(8.4 \pm 0.6) \cdot 10^{-6}$ the strength of this decay mode, as measured by the B_{GT}-value, is unexpectedly high. The result is discussed within a simple single-particle model and could be interpreted as a quasi-free decay of the ^{11}Be halo neutron into a single-proton state.

Keywords: beta decay, halo nucleus, ^{11}Be

1. Introduction

Beta-minus decay and proton emission take a nucleus in almost opposite directions on a nuclear chart, so β^--delayed proton emission (where beta decay feeds excited states that subsequently emit a proton) is forbidden in all but a few nuclei where it is heavily suppressed as the available energy is [1] $Q_{\beta p} = 782\,\mathrm{keV} - S_n$ where S_n is the neutron separation energy of the nucleus. We describe here an experiment to detect this decay mode from the one-neutron halo nucleus ^{11}Be that is believed to be the most favourable case [2, 3] due to the single-particle behaviour of halo nuclei [4, 5, 6] that may favour this decay mode and due also to the relatively long halflife that is caused by the normal beta-decay of ^{11}Be being hindered since a level inversion gives it a $1/2^+$ ground state rather than a $1/2^-$.

Beta-delayed particle emission is in general a prominent decay mode for nuclei close to the dripline, see [7, 8] for recent reviews. The energetically open channels for ^{11}Be are $\beta\alpha$, βt, βp and βn with corresponding Q-values of [9] 2845.2 ± 0.2 keV, 285.7 ± 0.2 keV, 280.7 ± 0.3 keV and 55.1 ± 0.5 keV. The low decay energy implies that the branching ratio for beta-delayed proton emission is low, typical estimates are slightly above 10^{-8} [3]. To detect the process experimentally, it is therefore essential to keep contaminants at a very low level.

The βp decay mode may be expected preferentially in one-neutron halo nuclei, partly due to the requirement of low neutron separation energy, partly due to the more pronounced single-particle behaviour of halo nuclei. Two-neutron halo nuclei are in a similar way candidates for beta-delayed deuteron emission, which has so far been observed only in the nuclei ^{6}He and ^{11}Li [7, 10]. For ^{11}Li the decay has a branching ratio of order 10^{-4}, the low value again caused by a small energy window, whereas cancellation effects reduces the branching ratio for ^{6}He down to the 10^{-6} level. It may be more useful to consider the standard measure for the strength of a decay, the reduced matrix element

*Corresponding author
Email address: kvr@phys.au.dk (K. Riisager)
[1] Present address: Institut für Kernphysik, Technische Universität Darmstadt, D–64289 Darmstadt, Germany

squared B_{GT}, that is found from the relation [7]

$$ft = \frac{K}{g_V^2 B_F + g_A^2 B_{GT}} \quad (1)$$

where f is the beta-decay phase space, $K/g_V^2 = 6144.2 \pm 1.6$ s and $g_A/g_V = -1.2694 \pm 0.0028$. Converting the observed spectra for beta-delayed deuteron emission from the two-neutron halo nuclei ^{6}He [11] and ^{11}Li [12] gives total B_{GT} values within the observed energy range of about 0.0016 and 0.75. (Note, however, that the ^{6}He decay to the ^{6}Li ground state has been described as an effective di-neutron to deuteron decay, it is a super-allowed transition with a B_{GT} of 4.7. This may be a reflection of a general trend for super-allowed decays to occur in very neutron-rich nuclei [13].) For comparison, the sum of B_{GT} for all currently known transitions in the ^{11}Be decay is 0.27.

2. The experiment

2.1. General remarks

The radioactive ^{11}Be nuclei were produced at the ISOLDE facility at CERN. Searching for protons with a kinetic energy of a few hundred keV with relative intensity 10^{-8} is challenging in a radioactive beam environment, so we instead detect the decay product, ^{10}Be with a halflife of $1.5 \cdot 10^6$ y, that exists only in minute quantities on earth. To reach the needed sensitivity we must employ state-of-the-art AMS. It is also crucial to limit the amount of contaminants in the samples so sample collection took place at ISOLDE's high-resolution mass separator. The resolution from the magnetic separation stage is supplemented by the electrostatic beam transport at ISOLDE similar to, but at lower resolution than, the separation stages in AMS facilities. A first attempt was made in 2001 and the results were published recently [3]. The signal was not sufficiently strong to be clearly separated from background and gave a βp branching ratio of $(2.5 \pm 2.5) \cdot 10^{-6}$, significantly above the published theoretical expectations. Due to improvements both in production of ^{11}Be and AMS detection of ^{10}Be, the current collection was performed in December 2012 and resulted in three samples.

2.2. Sample collection

The ^{11}Be activity was produced by bombarding a UC target with 1.4 GeV protons. The products were ionized in a laser ion source, which provided element selectivity, mass separated in the ISOLDE high-resolution separator, and guided through several collimators to the collection point where they were implanted at 60 keV in a small copper plate (15×20×2 mm). A high-purity coaxial Ge-detector placed 40 cm downstream behind a lead shielding monitored the collection rate. The Ge-detector was energy and efficiency calibrated with standard sources of ^{60}Co, ^{152}Eu and ^{228}Th. The main lines in the γ spectrum recorded during ^{11}Be collection are the 2124 keV line from the decay of ^{11}Be and the 511 keV line from positron annihilation. The overall efficiency at 2124 keV is found to be $(2.0 \pm 0.2) \cdot 10^{-5}$. A second line from the decay at 2895 keV was also used to check the overall amount of ^{11}Be. The two determinations gave about the same precision, the one from the 2124 keV line being dominated by systematic uncertainties in the efficiency and the one from the 2895 keV being dominated by statistical uncertainties, and were internally consistent leading to a final value for the amount of collected ^{11}Be in the main sample (S1) of $(1.447 \pm 0.055) \cdot 10^{12}$. This includes a correction for dead time of 2.8%, determined from the ratio of accepted to total number of triggers.

As cross-checks two other samples were collected: sample S2 at the mass position of ^{11}Li (0.02 mass units heavier than ^{11}Be) where an upper limit of $3 \cdot 10^6$ could be determined for the number of atoms collected (corresponding to a ^{11}Li yield below 625/s which is reasonable) and, for one second only, sample S3 at the ^{10}Be mass position where an estimate of the current of 3.5 pA (uncertain by a factor two) converts into $2.2 \cdot 10^7$ atoms. According to SRIM calculations [14] about 6% of all Be ions implanted in Cu at 60 keV energy will backscatter out of the sample. Most of the backscattered ions are expected to remain close to the sample so γ-rays from their decays will be seen as well, although the decay products are not retained in the sample. This gives a correction which we estimate to be $4 \pm 4\%$.

2.3. Accelerator mass spectrometry

The ^{10}Be accelerator mass spectrometry (AMS) measurements were performed at the Vienna Environmental Research Accelerator (VERA) at the University of Vienna. VERA is a dedicated AMS facility based on a NEC 3 MV pelletron tandem accelerator. A new scheme for ^{10}Be using a passive foil absorber in front of a gas ionization chamber detector was employed. In this way the detection efficiency for ^{10}Be atoms is increased significantly.

According to TRIM simulations [14] the maximum implantation depth of ^{11}Be in our copper plate catcher was below 1 μm. To reduce the amount of material to be dissolved only the surface layer of each irradiated copper plate was leached in nitric acid. A second leaching was performed to verify the blank level of the irradiated copper plate. The second leaching of sample S3 did not produce enough BeO for a measurement. For samples S1 and S2 the values of the second leachings were consistent with a blank sample. This shows that the material was sitting in the surface, as expected for an implanted sample, and not due to a bulk contamination. An amount of 359 μg (uncertainty 3%) ^{9}Be carrier was added to the solution to reach a ^{10}Be/^{9}Be isotopic ratio in the range of 10^{-16}–10^{-11}. In the next step the solution was treated with ammonium hydroxide to precipitate the beryllium as beryllium hydroxide ($Be(OH)_2$). The dissolved copper remains in the solution in this step. The beryllium hydroxide was dried out by heating in an oven at 900°C for at least 8 hours forming beryllium oxide (BeO). The BeO powder was mixed 1:1 with high purity copper powder and pressed into sample holders and mounted together with standard and blank material in a MC-SNICS type Cesium sputter ion source. Blank is the pure phenakite material directly pressed into a sample holder. A separate sample, S-blank, went through the chemistry preparation to check for the amount of ^{10}Be introduced during the chemical sample preparation. BeO^- was extracted from the ion source and stripped in the terminal of the tandem accelerator to Be^{2+}, resulting in a total ion energy of 2.4 MeV. After further mass separation by a sector magnetic analyzer and an electrostatic analyzer the remaining particles are sent to a gas ionization chamber detector with a two-split anode for particle identification. A silicon nitride foil stack as a passive absorber was installed in front of the detector. This foil stack prevents the isobaric background ^{10}B from entering into the detector: The energy loss of boron in the foil stack is slightly larger compared to beryllium. By selecting the right foil thickness and carefully tuning the particle energy the boron ions are stopped in the foil stack whereas the beryllium ions can enter the detector.

The final results are given in table 1. The amount of atoms in sample S3 agrees with the estimation from the implantation current. The number for sample S2 is consistent with the lack of observed γ-rays from the decay of ^{11}Li. The number for sample S1, the ^{11}Be sample, is $(1.170 \pm 0.047) \cdot 10^7$.

Table 1: Results of the AMS measurement. S1 to S3 denote the irradiated samples. 1st and 2nd correspond to the first or second leaching. Blank and S-blank are control samples without activity.

Sample	^{10}Be/^{9}Be ratio	^{10}Be atoms
S1-1st	$(4.87 \pm 0.13) \cdot 10^{-13}$	$(1.17 \pm 0.05) \cdot 10^7$
S1-2nd	$(1.26 \pm 0.56) \cdot 10^{-15}$	$(3.03 \pm 1.35) \cdot 10^4$
S2-1st	$(3.10 \pm 0.94) \cdot 10^{-15}$	$(7.45 \pm 2.27) \cdot 10^4$
S2-2nd	$(4.4 \pm 3.1) \cdot 10^{-16}$	$(1.06 \pm 0.75) \cdot 10^4$
S3-1st	$(1.54 \pm 0.03) \cdot 10^{-12}$	$(3.70 \pm 0.13) \cdot 10^7$
S-blank	$(4.9 \pm 3.4) \cdot 10^{-16}$	$(1.18 \pm 0.82) \cdot 10^4$
blank	$(1.3 \pm 1.3) \cdot 10^{-16}$	$(3.12 \pm 3.12) \cdot 10^3$



Figure 1: Mass scan of the ISOLDE high-resolution separator across the ^{11}Be position. The beta activity measured is shown versus the mass with positions indicated for ^{11}Be and the possible contaminant ^{11}Li. The horizontal line marks the detection limit of 0.2/s.

2.4. Possible contaminants

Contaminations in our sample might arise due to tails of the neighbouring activities ^{10}Be or ^{11}Li, whose decay also produces ^{10}Be. Both possibilities are ruled out by the low recorded number of atoms for the ^{11}Li sample (S2). The ISOLDE mass separator profile was found by measuring the beta activity as the mass settings were changed around the nominal ^{11}Be mass, see figure 1. The release function of this specific target and ion source combination was measured first, which allows to combine measurements with different collection times relative to

proton impact on target. In this way the sensitivity was increased and the activity could be followed down to the 10^{-5} level that occured at a mass difference of 0.05 mass units. The only remaining way for ^{10}Be to appear on the ^{11}Be position is as the molecule ^{10}Be^{1}H, but this molecule is unlikely to be formed in the target and to survive through the laser ion source since its ionization energy of 8.22 eV [15] is much higher than its dissociation energy of 3.26 eV. Nevertheless, we have re-checked the data from an earlier experiment on ^{12}Be [16] and were able to put limits on the amount of ^{11}Be^{1}H (from the $\beta\alpha$ branch) that would correspond in our current case to a ^{10}Be^{1}H intensity less than $2 \cdot 10^{-6}$ of ^{11}Be. Our conditions should be better, partly due to higher laser ionization power, partly due to the beam passing through a gas-filled RFQ cooler, both effects that would enhance molecular breakup. We therefore conclude that we have observed the ^{11}Be(βp) decay via detection of the final nucleus ^{10}Be. The observed intensity converts to a branching ratio of $(8.4 \pm 0.6) \cdot 10^{-6}$.

3. Discussion

The experimentally found branching ratio is surprisingly large, but consistent with the outcome of the first experiment. If the strength in ^{11}Be(βp) was as broadly distributed as in ^{11}Li(βd) we would expect the B_{GT} within the Q-window to be less than 0.1, which would not be sufficient to explain the decay rate. We therefore turn to a simple model for the decay along the lines of the direct decay calculations in [2, 17], details of the calculations are reported elsewhere [18].

The basic assumption is that the beta decay proceeds as an essentially detached decay of the halo neutron into a proton. The initial and final state wavefunctions are calculated as single-particle states in square-well or Woods-Saxon potentials with the final state spectrum discretized by imposing a large confining radius at 1000 fm. The overlap of the wavefunctions gives the beta strength B_{GT} and the decay rate is found from equation (1). The final total branching ratio for beta-delayed proton emission depends strongly on the strength of the potential between the final state proton and ^{10}Be. For most potential strengths the branching ratio will indeed be a few times 10^{-8}, as in other calculations, but in a limited range the beta strength will be concentrated within the Q-window. Effectively, in this range the proton formed in the decay interacts strongly with the remaining ^{10}Be and forms a resonance-like structure; as a consequence it emerges with a quite well defined energy. The branching ratios obtained for this set of parameters are shown in figure 2 as a function of the energy of the resonance.



Figure 2: The calculated branching ratio for decay into p+^{10}Be is shown as a function of the centre-of-mass energy of the resonance. The stars mark results of calculations with square well potentials, the filled circles are results from Woods-Saxon potentials. The curves arise from integrating the R-matrix expression in equation (2) for different widths for other decay channels. The horizontal band indicates the experimentally found branching ratio.

The simple model neglects isospin. The lowest $T = 3/2$ states are situated slightly more than 1 MeV above the $Q_{\beta p}$ window. They are members of isospin multiplets that include the ^{11}Be ground state and first excited state neutron halos. The data indicate [19] that the intermediate states ($|T_z|$ of 1/2) in these multiplets have good total isospin rather than a composition with just one proton (or neutron) plus core. We therefore expect that realistic final state wave functions in our case, with $T = 1/2$, also should have good isospin. Standard isospin coupling then predicts that the state should be proton plus ^{10}Be with weight 2/3 and neutron plus ^{10}B(T=1) with weight 1/3. Our calculated decay probabilities must therefore be corrected by a factor 2/3. A further reduction factor about 0.7 is due to the initial ^{11}Be wavefunction containing several configurations [20]. The overall scaling factor on the theory, included in figure 2, is therefore about 0.5.

Could this be an established resonance in ^{11}B ?

The known states [21] in this region mainly couple to the α-particle channel (with partial widths around 100 keV) and only one, a state at 11450 ± 17 keV, may have spin-parity that allows emission of an s-wave proton — the others will have angular momentum barriers that will suppress proton emission. Decays through levels that have other sizeable decay channels (α emission or, for very narrow levels, γ emission; in principle triton emission could also occur) would therefore only contribute to the proton channel with probability Γ_p/Γ_{tot}. Since Γ_γ for the $1/2^+$, $T = 3/2$ state at 12.55 MeV (that apart from isospin should be similar in structure to our state) is about 10 eV [21], and even a small admixture into our state of other $1/2^+$ levels is likely to give a Γ_α at least of the same magnitude, we shall assume the width for other decay channels $\Gamma_b = \Gamma_\gamma + \Gamma_\alpha$ to be larger than 0.01 keV.

To take these effects into account calculations were also made within the R-matrix approach [22], but in a simplified version where e.g. other decay channels are approximated as having a constant width Γ_α over the energy window, see [18] for details. Converting the decay rate into a differential branching ratio gives the following expression:

$$\frac{\mathrm{d}b}{\mathrm{d}E} = t_{1/2}\frac{g_A^2}{K}\frac{B_{GT}\Gamma_p/2\pi}{(E_{res}-E)^2+\Gamma_{tot}^2/4}f(Q-E)\,, \quad (2)$$

where $\Gamma_{tot} = \Gamma_b + \Gamma_p$, $\Gamma_p = 2P\gamma^2$, P is the standard (energy-dependent) penetrability factor and γ^2 the maximal reduced width. Integration over the Q-window gives the total branching ratios shown in figure 2 as a function of resonance position E_{res} for different values of Γ_b. The branching ratios agree well with the ones from the simple model. It is clear that all known levels are too wide to fit and that a Γ_b above 0.01 keV gives a lower limit on the B_{GT} of about 0.3 with an upper limit given by the theoretical maximum of 3.

4. Conclusion

We have observed beta-delayed proton emission for the first time in a neutron-rich nucleus. The unexpectedly high decay rate can only be understood within current theory if the decay proceeds through a new single-particle resonance in ^{11}B that is strongly fed in beta-decay. The B_{GT}-value could be as high as that of a free neutron decay. A natural interpretation would be peripheral beta decay of the halo neutron in ^{11}Be into a single-proton state. This appears to be a simpler process than the βd decays of the two-neutron halo nuclei ^{6}He and ^{11}Li. Although the halo structure must be important for the βp decay mode, the large value of B_{GT} may be related to large values found in other (non-halo) near-dripline nuclei [13] and point to a more widespread change of beta-decay patterns at least in light nuclei in line with some predictions [23].

Acknowledgements

We thank the ISOLDE group for the successful operation of the HRS separator at very high resolution and Aksel Jensen for discussions on the theoretical interpretation. We acknowledge support from the European Union Seventh Framework through ENSAR (contract no. 262010), from Austrian Science Fund (FWF) P22164-N20, from Spanish MINECO through projects FPA2010-17142 and FPA2012-32443, and CPAN Consolider CSD-2007-00042.

References

References

[1] B. Jonson and K. Riisager, Nucl. Phys. A **693** (2001) 77.
[2] D. Baye and E.M. Tursonov, Phys. Lett. B **696** (2011) 464.
[3] M.J.G. Borge et al., J. Phys. G **40** (2013) 035109.
[4] A.S. Jensen, K. Riisager, D.V. Fedorov and E. Garrido, Rev. Mod. Phys. **76** (2004) 215.
[5] I. Tanihata, H. Savajols and R. Kanungo, Prog. Part. Nucl. Phys. **68** (2013) 215.
[6] K. Riisager, Physica Scripta T**152** (2013) 014001.
[7] M. Pfützner, M. Karny, L.V. Grigorenko and K. Riisager, Rev. Mod. Phys. **84** (2012) 567.
[8] B. Blank B and M.J.G. Borge, Prog. Part. Nucl. Phys. **60** (2008) 403.
[9] M. Wang et al., Chinese Phys. C **36** (2012) 1603.
[10] T. Nilsson, G. Nyman and K. Riisager, Hyperfine Int. **129** (2000) 67.
[11] D. Anthony et al., Phys. Rev. C **65** (2002) 034310.
[12] R. Raabe et al., Phys. Rev. Lett. **101** (2008) 212501.
[13] M.J.G. Borge et al., Z. Phys. A **340** (1991) 255.
[14] J.F. Ziegler, Particle interactions with matter. http://www.srim.org (Jan. 20, 2014)
[15] S. Bubin and L. Adamowicz, J.Chem.Phys. **126** (2007) 214305.
[16] C.Aa. Diget et al., Nucl. Phys. A **760** (2005) 3.
[17] M.V. Zhukov, B.V. Danilin, L.V. Grigorenko and J.S. Vaagen, Phys. Rev. C **52** (1995) 2461.
[18] K. Riisager, submitted to Nucl. Phys. A (2014), arXiv:1312.0479
[19] B. Jonson and K. Riisager, Phil. Trans. R. Soc. Lond. A **356** (1998) 2063.

[20] K.T. Schmitt et al., Phys. Rev. Lett. **108** (2012) 192701.
[21] J.H. Kelley et al., Nucl. Phys. A **880** (2012) 88.
[22] F.C. Barker and E.K. Warburton, Nucl. Phys. A **487** (1988) 269.
[23] H. Sagawa, I. Hamamoto and M. Ishihara, Phys. Lett. B **303** (1993) 215.